

AN ANALYSIS OF DDT RESIDUES IN MILK SAMPLES OF BUFFALO FROM KANPUR DISTRICT (U.P.), INDIA

ALOK KUMAR PANDEY*

ABSTRACT

In all 62 milk samples from buffalo of 7 regions of Kanpur district were analysed. The results (Table-1) revealed that 37.68 percent samples were contaminated with DDT in the range of 0.003 to 1.749 ppm, of which 4.52 percent samples exceeded the maximum permissible limit of 1.25 ppm. More than 50 percent of the buffalo milk samples (54.83%) were contaminated with DDT in the range of 0.014 to 1.749 ppm. From these samples 23.53 percent samples exceeded the permissible level of 1.25 ppm.

KEYWORDS: Milk Sample, Buffalo, DDT.

INTRODUCTION

Pesticides have come in for severe criticism at various forums in the world and India is no exception, since, the publication of Rachel Carson's "Silent Sprint" in 1962 with its emotional denunciation of chemical pesticides. Pesticides are an essential part of production technology that makes it possible for the farmers to meet the demands of our growth population. Without pesticides, crop pest would make monoculture impossible and this would also make scientific farming impracticable. Plant diseases and insect pests could spell disaster in any given year, were it not for the protection of agriculture. If we do not use pesticides, it is estimated that crop losses would be high as 70-100 percent on an average and in some cases, pest damage or competition would make is pointless even to plant the crop. Substantial gains from the use of pesticides in agriculture as well as public health programme have been acknowledge by all.

None the less, the harmful residual that remain on the crop, especially edible portion has cause great concern to everyone. Besides the contamination of feed and food commodities with pesticide, there is a growing, risk, that pesticides particularly which persist and then types that get translocated through air, water and soil system from the point of initial application, thus endanger the normal functioning of non target organisms. Bioaccumulation of these pesticides in food commodities and other environmental components is an important area, which needs immediate attention. About 120 pesticides are registered for use in India of which two insecticides DDT and BHC most common, because of their efficacy as broad spectrum, to insect pests in agriculture and in public health, occupied with reasonable low prices compared to other organochlorine, organophosphorus, carbamate and pyrethroid insecticides.

*Department of Zoology, D.A.V. P.G. College, Civil Lines, Kanpur.

Correspondence E-mail Id: editor@eurekajournals.com

MATERIAL AND METHOD

The present investigation was undertaken with the objective of monitoring DDT and BHC residues in market/ field samples of various commodities like okra, milk and water from Kanpur Districts of U.P. State (India). The analysis of DDT and BHC residue estimation was done as per the procedure described in ISI bulletin, AFDC-56 (2326) P, (1982) and AFDC 56 (2332) PC 1982 with little modifications. The materials used and methods followed are described below. Chemicals required of DDT and BHC residue estimation. Sodium Chloride, Anhydrous sodium sulphate, Acetone (redistilled), Methyl cyanide (redistilled), n-hexane (redistilled), Petroleum ether (redistilled), Sulphuric acid (sp. gr. 1.84), m,p'; and p,p' and p,p' – DDT isomers, Alpha, Beta and Gamma HCH. Animal milk samples were homogenized by vigorous shaking and reduced to 100 ml and 5 ml respectively. Animal milk samples measuring 100 ml and 5 ml respectively were taken in blender, extracted in acetone + hexane mixture (1:1). Hexane extract (50ml) was transferred in separatory funnel and about 100 ml sulphuric acid was added drop wise till hexane layer got clear. The lower phase of sulphuric acid was discarded. The upper phase was washed with distilled water till it became neutral, dried over anhydrous sodium sulphate and concentrated to 10ml.

RESULT AND DISCUSSION

In all 62 milk samples from buffalo of 7 regions of Kanpur district were analysed. The results

(Table-1) revealed that 37.68 percent samples were contaminated with DDT in the range of 0.003 to 1.749 ppm, of which 4.52 percent samples exceeded the maximum permissible limit of 1.25 ppm. More than 50 percent of the buffalo milk samples (54.83%) were contaminated with DDT in the range of 0.014 to 1.749 ppm. From these samples 23.53 percent samples exceeded the permissible level of 1.25 ppm. Such milk samples were unfit for direct consumption. Kapoor et al. (1980) reported that buffalo milk samples collected from two regions were contaminated with DDT and there was direct correlation found between concentrations of DDT in the milk and DDT use in malaria control. In Kanpur cottonseed is directly fed to the mulching buffaloes for improving quality and quantity of milk. Contaminated cottonseed may help in contamination of milk with DDT. Similar results are also found by Dhaliwal, *et. al.* (1977) DDT residues in milk samples from Ludhiana and surrounding areas, Khan, *et. al.* (2013), Mathiyazhagan, N., & Natarajan, D. (2013). Phytoremediation efficiency of edible and economical crops on waste dumps of bauxite mines, Salem district, Tamil Nadu. On a sustainable future of the Earth natural resources, Nassar, *et. al.* (2018), Ozemen-M, *et. al.* (2008), Ochiai, M. and Hanya, T. (1976). Alpha and gamma BHC in Tamgava river water, Japan, Rozpara E, *et. al.* (2016), Sarwar M, Salman M (2015), Senthil-Nathan S (2015), Shahab-Ghayoor H, Saeidi K (2015), Szolyga B, *et. al.* (2014) and Tak JH *et. al.* (2016).

Table 1.DDT residues in milk samples from Kanpur district (U.P.), India

S. No.	Commodity	No. of samples analysed	No. of samples containing DDT residues				Residue level	
			BDL	No. contaminated sample	1.25ppm	1.25 ppm	Min.	Max.
1	Buffalo milk	62	28 (41.17)	34	26 (94.59)* (76.47)*	(5.40)* (23.52)*	0.014	1.749

* Based on 5 observations

** Figures in brackets are $\sqrt{n+1}$ transformations

*** Significant

Fig.In Parentheses and Transformed value.

REFERENCES

- [1]. Dhaliwal, G.S. and Kalra, R.K. (1977) DDT residues in milk samples from Ludhiana and surrounding areas. Indian J. ecol., 4(1): 13-22.
- [2]. Khan, H.A., Akram, W. Shad, S.A. & Lee, J.J. (2013). Insecticide mixtures could enhance the toxicity of insecticides in a resistant dairy population of *Musca domestica* L. *PloS One*, 8(4), 609-629.
- [3]. Mathiyazhagan, N., & Natarajan, D. (2013). Phytoremediation efficiency of edible and economical crops on waste dumps of bauxite mines, Salem district, Tamil Nadu. On a sustainable future of the Earth natural resources. Springer earth system science. DOI <https://doi.org/10.1007/978-3-642-32917-3-31>.
- [4]. Nassar, M.I., Ghazawy, N.A., Torkey, H.M., & Rawy, S.M. (2018). Assessment of biorational botanical extracts on the desert locust *Schistocerca gregaria* Forskal (Orthoptera: Acrididae). *Entomol Appl Sci Lett*, 5(2), 42-54.
- [5]. Ozemen-M, Ayaas -Z, Gungardon-A (2008). Given the effects of environmental pollution and different biotic factors or some important biochemical marker as enzyme.
- [6]. Ochiai, M. and Hanya, T. (1976). Alpha and gamma BHC in Tamgava river water, Japan (September, 1968- September, 1969). *Environ., Pollut.*, 11(3) 161-166.
- [7]. Rozpara E, Morgas H, Filipczak J, Meszka B, Holdaj M. Labanowska BH, Sekrecka M, Sobiczewski P, Lisek J, Danelski W (2016) Metodyka produkcji owocow aronii metoda ekologiczna (wydanie pierwsze), Inhort, Skierniewice, p 27.
- [8]. Sarwar M, Salman M (2015) Toxicity of oils formulation as a new useful tool in crop protection for insect pests control. *Int J Chem Biomol Sci* 1:297-302.
- [9]. Senthil-Nathan S (2015) A review of biopesticides and their mode of action against insect pests. In: Thangavel P, Sridevi G (eds) *Environmental Sustainability*. Springer, Delhi, pp 49-63.
- [10]. Shahab-Ghayoor H, Saeidi K (2015) Antifeedant activities of essential oils of *Satureja hortensis* and *Fumaria parviflora* against Indian Meal Moth *Plodia interpunctella* Hubner (Lepidoptera; Pyralidae). *Entomol Ornithol Herpetol* 4:1-4, <https://doi.org/10.4172/2161-0983.1000154>.
- [11]. Szolyga B, Gnilka R, Szumny Szczepanik M (2014) Chemical composition and insecticidal activity of *Thuja occidentalis* and *Tanacetum vulgare* essential oils against larvae of the lesser mealworm, *Alphitobius diaperinus*. *Entomol Exp Appl*

- 151: 1-10. <https://doi.org/10.1111/eea.12166>.
- [12]. Tak JH, Jovel E, Isman MB (2016) Contact, fumigant, and cytotoxic activities of thyme and lemongrass essential oils against larvae and an ovarian cell line of the cabbage looper, *Trichoplusia ni*. *J Pest Sci* 89:183-193.